



W. Garrett Owen¹



Josh Henry²



Brian E. Whipker²

Volume 2 Number 13 August 2019

Nutritional Monitoring Series

Poinsettia

(*Euphorbia pulcherrima*)

Poinsettias require medium to high fertility levels between 150 and 300 ppm N, with the rate adjusted to take into account plant vigor. Optimal substrate pH values for poinsettia range from 5.5 to 6.5. If the substrate pH drifts lower than 5.5, plants will become stunted and color delayed. Substrate pH values above 6.5 inhibit Fe availability and induce interveinal chlorosis and stunted plant growth.



Funding Generations of Progress
Through Research and Scholarships



Project Sponsor



Figure 1. Low substrate pH will result in stunted poinsettia (*Euphorbia pulcherrima*) plant growth. Photo by: Brian Whipker.

Target Nutrition Parameters

pH Category II - IV:
5.5 to 6.5

Fertility Category:
Medium to High
150 to 300 ppm N

EC Category B - C:
1:2 Extraction:
0.6 to 1.5 mS/cm

SME:
1.3 to 3.3 mS/cm

PourThru:
2.0 to 4.3 mS/cm

Poinsettia

www.fertdirtandsquirt.com

¹Michigan State University
wgowen@msu.edu

²NC State University
bwhipker@ncsu.edu



Figure 2. Lower leaves of poinsettia (*Euphorbia pulcherrima*) exhibiting lower leaf interveinal chlorosis (yellowing) due to a low substrate pH-induced magnesium (Mg) deficiency. Photo by: W. Garrett Owen.



Figure 3. Substrate pH >6.5 can inhibit iron (Fe) uptake causing poinsettia (*Euphorbia pulcherrima*) leaves to become Fe-deficient and exhibit interveinal chlorosis (yellowing). Leaf tissue nutrient analysis determined the mature leaves under the bracts to exhibited Fe-deficiency (82.8 ppm Fe). Photo by: W. Garrett Owen.



Figure 4. As a result of high substrate pH, iron (Fe)-deficient poinsettia (*Euphorbia pulcherrima*) exhibit intensified interveinal chlorosis where recently matured leaves become entirely chlorotic (yellow). Photo by: Brian Whipker.

Fertility Management of Poinsettia

Poinsettias should be grown with a substrate pH range of 5.5 to 6.5. Use recommended 1:2 Extraction, SME, or PourThru methods to determine and monitor substrate pH and soluble salts [referred to as electrical conductivity (EC)] values. Additionally, conduct routine foliar analysis tests to monitor crop nutrient status. Tissue nutrient levels found in healthy, newly expanded poinsettia leaves and critical tissue values are provided in Table 1, which can help in diagnosing suspected nutrient disorders. Monitoring substrate pH and nutrient status will enable growers to avoid pH induced nutritional disorders.

Substrate pH below 5.5 will result in stunted plants (Fig. 1) and delay color development without any lower leaf bronzing or black spotting (Whipker, 2014). Corrective procedures for low substrate pH should begin at 5.4. Substrate pH below 5.4 to 5.6 may inhibit magnesium (Mg) uptake causing lower or older leaves to become Mg-deficient and exhibit interveinal chlorosis (yellowing; Fig. 2). Monthly applications of supplemental Mg in the form of magnesium sulfate (MgSO₄; Epsom salts) at a rate of 8 oz./100 gal. of water in areas with naturally occurring Mg in the water supply or 16 oz./100 gal. of water in areas lacking Mg in the irrigation water will prevent Mg deficiency and symptomology development.

High substrate pH above 6.5 can inhibit Fe uptake causing newly developed and recently matured leaves to become Fe-deficient (Fig. 3). Symptoms of Fe deficiency initially appear as slight chlorosis, intensifying to interveinal chlorosis (Fig. 4), and in severe instances, plants will become completely chlorotic (Fig. 5). Plant growth may also



Figure 5. As a result of high substrate pH, iron (Fe) deficiency of poinsettia (*Euphorbia pulcherrima*) intensifies causing the plant to become completely chlorotic (yellow). In this photo, leaf tissue nutrient analysis determined Fe-deficiency (47.7 ppm Fe). Photo by: W. Garrett Owen.



Figure 6. High substrate pH above 6.5 can result in stunted poinsettia (*Euphorbia pulcherrima*) plant growth. Photo by: Brian Whipker.



Figure 7. Providing insufficient fertility [low electrical conductivity (EC)] during poinsettia (*Euphorbia pulcherrima*) production can result in stunted plant growth. Photo by: Brian Whipker.

become stunted (Fig. 6). Corrective procedures for high substrate pH should begin within the range of 6.2 to 6.4.

For more information about pH disorders, refer to [e-GRO Alert 3-59: Poinsettia Fertilization: pH Disorders](#).

During poinsettia production, maintain a medium to high level of fertility at 150 to 300 ppm N, but modify for plant vigor. Insufficient fertility levels (low EC) will result in stunted plant growth (Fig. 7). Overfertilization (high EC) results in excessive growth (Fig. 8) and marginal leaf necrosis [death; (Fig. 9)]. If EC levels begin to increase, stop fertilization and simply apply a few clear water irrigations. This corrective procedure will aid in lowering substrate EC. Prior to resuming fertilization, perform a 1:2 Extraction, SME, or PourThru to determine if the EC levels have dropped back into the acceptable range. If EC values become excessive, leach the substrate with clear irrigation water twice before providing fertility. It is best to monitor the crop to avoid excessive EC values than to waste fertilizer by leaching it from the pots.

Summary

Providing medium to high fertility at 150 to 300 ppm N and maintaining a substrate pH of 5.5 to 6.5 will help prevent most nutritional disorders in poinsettia.

Literature Cited

Dole, J.M. and H.F. Wilkins. 2005. Floriculture: Principles and species. 2nd ed. Pearson Education, Inc., Upper Saddle River, NJ.



Figure 8. Excessive fertility [high electrical conductivity (EC)] during poinsettia (*Euphorbia pulcherrima*) production can result in excessive plant growth. Photo by: Brian Whipker.



Figure 9. Excessive fertility [high electrical conductivity (EC)] during poinsettia (*Euphorbia pulcherrima*) production can result in marginal leaf necrosis. Photo by: Brian Whipker.

Literature Cited (continued)

Ecke, P. III, J.E. Faust, J. Williams, and A. Higgins. 2004. The Ecke poinsettia manual. Ball Pub., West Chicago, IL.

Whipker, B.E. 2014. Poinsettia fertilization: pH disorders. 3:1-5.

Table 1. Critical, recommended, and toxic leaf tissue nutrient concentrations for poinsettia (*Euphorbia pulcherrima*).

Element		Critical ¹	Recommended ^{1, 2}	Toxic ^{1, 2}
Nitrogen (N)	(%)	3.5	4.0 - 6.0	7.0 - 7.3
Phosphorus (P)		0.2	0.3 - 0.6	0.8 - 0.9
Potassium (K)		1.0	1.5 - 3.5	4.0
Calcium (Ca)		0.5	0.7 - 1.8	-
Magnesium (Mg)		0.2	0.3 - 1.0	-
Sulfur (S)		0.05	0.1 - 0.3	-
Sodium (Na)		-	0 - 0.4	0.5
Chloride (Cl)		-	0 - 1.5	3.0
Iron (Fe)	(ppm)	50	100 - 300	-
Manganese (Mn)		40	60 - 300	650
Zinc (Zn)		20	25 - 60	-
Copper (Cu)		1	2 - 10	-
Boron (B)		15	25 - 75	100
Molybdenum (Mo)		0.5	1 - 5	-
Fluoride (F)		-	0 - 4	5
Lithium (Li)		-	0 - 15	20

Sources: ¹ Ecke et al. (2004); ² Dole and Wilkins (2005)

Corrective Procedures for Modifying Substrate pH and Electrical Conductivity (EC)

When the pH or substrate electrical conductivity (EC) drifts into unwanted territory, adjustments must be made. Below are the standard corrective procedures used to modify the substrate pH and EC for greenhouse grown crops in soilless substrates.

1. Low Substrate pH Correction

When Fe and Mn toxicity becomes a problem, adjust (raising) substrate pH to the recommended pH range. Corrective procedures to raise low pH levels are listed below. Switching to a basic fertilizer when the substrate pH is nearing the lower limit will help stabilize the pH. If the pH is below the recommended range, then corrective procedures will need to be implemented. Flowable lime is one option. Using a rate of 2 quarts per 100 gallons of water will typically increase the substrate pH by roughly 0.5 pH units. Two quarts can be used through an injector. Additional applications can be made if needed. Potassium bicarbonate (KHCO_3) can also be applied. A rate of 2 pounds per 100 gallons of water will increase the substrate pH by roughly 0.8 pH units. This treatment will also provide excessive potassium (K) and cause a spike in the substrate EC. A leaching irrigation with clear water is required the following day to restore the nutrient balance (the ratio of K:Ca:Mg) and lower the EC. As always, remember to recheck your substrate pH to determine if reapplications are needed.

pH Adjustment Recommendations

Flowable Lime

- Use 1 to 2 quarts per 100 gallons of water.
Rinse foliage.
- Avoid damage to your injector by using rates of 2 quarts per 100 gallons of water, or less.
- Can split applications.

Hydrated Lime

- Mix 1 pound in 3 to 5 gallons of WARM water. Mix twice. Let settle. Decant liquid and apply through injector at 1:15.
- Caustic (rinse foliage ASAP and avoid skin contact)

Potassium Bicarbonate (KHCO_3)

- Use 2 pounds per 100 gallons of water
- Rinse foliage immediately.
- Provides 933 ppm K.
- Leach heavily the following day with a complete fertilizer to reduce substrate EC and restore nutrient balance.
- Rates greater than 2 pounds per 100 gallons of water can cause phytotoxicity!

2. High Substrate pH Correction

The target pH for many species is between 5.8 and 6.2. Higher pH values will result in Fe deficiency and lead to the development of interveinal chlorosis on the upper leaves. Check the substrate pH to determine if it is too high. Be careful when lowering the substrate pH, because going too low can be much more problematic and difficult to deal with.

Acid-based Fertilizer

If the substrate pH is just beginning to increase, then first consider switching to an acidic-based fertilizer. These ammoniacal-nitrogen (N) based fertilizers are naturally acidic and plant nitrogen uptake will help moderate the substrate pH over a week or two.

Acid Water Drench

Some growers use this intermediate correction if pH levels are not excessively high and a quick lower of the substrate pH is desired. Use sulfuric acid to acidify your irrigation water to a pH 4.0 to 4.5. Apply this acid water as a substrate drench providing 5 to 10% excessive leaching of the substrate. Rinse the foliage to avoid phytotoxicity. Results should be visible within 5 days. Retest the substrate pH and repeat if needed.

Iron Drench

If the levels are excessively high, then an Fe chelate application can be made to the substrate.

Below are the options.

Iron Chelate Drench (options)

- Iron-EDDHA: mix 5 ounces in 100 gallons of water
- Iron-DTPA: mix 5 ounces in 100 gallons of water
- Iron sulfate: mix 4-8 ounces in 100 gallons of water
- Apply as a substrate drench with sufficient volume to leach the pot.
- Rinse foliage immediately.
- Avoid use on iron efficient plants (geraniums).

3. Low EC Correction

If low EC problems occur, increase the fertilization rate to 300 ppm N for a few applications before returning to the recommend fertilization rate for the crop.

4. High EC Correction

Excessively high fertilization rates will result in a marginal leaf burn. Check the substrate EC to confirm your diagnosis. Values greater than 6.0 mS/cm based on the PourThru sampling method can be problematic for many plants.

Switch to Clear Water Irrigations

If the substrate EC is just beginning to increase over time, then leach with a few clear water irrigations to lower EC levels by flushing out the salts.

Clear Water Leaching

If the EC values are excessively high, leach the substrate twice with back-to-back clear water irrigations. Then allow the substrate to dry down normally before retesting the EC. If EC levels are still too high, repeat the double leach. Once the substrate EC is back within the normal range, use a balanced fertilizer at a rate of 150 to 200 ppm N.



e-GROAlert

www.e-gro.org

CONTRIBUTORS

Dr. Nora Cattlin
Floriculture Specialist
Cornell Cooperative Extension
Suffolk County
nora.cattlin@cornell.edu

Dr. Chris Currey
Assistant Professor of Floriculture
Iowa State University
ccurrey@iastate.edu

Dr. Ryan Dickson
Greenhouse Horticulture and
Controlled-Environment Agriculture
University of Arkansas
ryand@uark.edu

Nick Flax
Commercial Horticulture Educator
Penn State Extension
nzf123@psu.edu

Thomas Ford
Commercial Horticulture Educator
Penn State Extension
tf2@psu.edu

Dan Gilrein
Entomology Specialist
Cornell Cooperative Extension
Suffolk County
dgg1@cornell.edu

Dr. Joyce Latimer
Floriculture Extension & Research
Virginia Tech
jlatime@vt.edu

Heidi Lindberg
Floriculture Extension Educator
Michigan State University
wolleage@anr.msu.edu

Dr. Roberto Lopez
Floriculture Extension & Research
Michigan State University
rllopez@msu.edu

Dr. Neil Mattson
Greenhouse Research & Extension
Cornell University
neil.mattson@cornell.edu

Dr. W. Garrett Owen
Floriculture Outreach Specialist
Michigan State University
wgowen@msu.edu

Dr. Rosa E. Raudales
Greenhouse Extension Specialist
University of Connecticut
rosa.raudales@uconn.edu

Dr. Beth Scheckelhoff
Extension Educator - Greenhouse Systems
The Ohio State University
scheckelhoff.11@osu.edu

Dr. Paul Thomas
Floriculture Extension & Research
University of Georgia
pthomas@uga.edu

Dr. Ariana Torres-Bravo
Horticulture / Ag. Economics
Purdue University
torres2@purdue.edu


Dr. Brian Whipker
Floriculture Extension & Research
NC State University
bwhipker@ncsu.edu

Dr. Jean Williams-Woodward
Ornamental Extension Plant Pathologist
University of Georgia
jwoodwar@uga.edu

Copyright © 2019

Where trade names, proprietary products, or specific equipment are listed, no discrimination is intended and no endorsement, guarantee or warranty is implied by the authors, universities or associations.

Cooperating Universities



Cornell University IOWA STATE UNIVERSITY



University of
New Hampshire
Cooperative Extension



PennState Extension



VIRGINIA
TECH

MICHIGAN STATE
UNIVERSITY

UConn

PURDUE
UNIVERSITY



The University of Georgia



THE OHIO STATE
UNIVERSITY

NC STATE
UNIVERSITY



DIVISION OF AGRICULTURE
RESEARCH & EXTENSION
University of Arkansas System

In cooperation with our local and state greenhouse organizations

MAUMEE VALLEY GROWERS
Choose the Very Best.



Metro Detroit Flower Growers Association



CONNECTICUT
GREENHOUSE
GROWERS
ASSOCIATION



Indiana
FLOWER
GROWERS
Association



Michigan Floriculture Growers Council

