



W. Garrett Owen<sup>1</sup>



Josh Henry<sup>2</sup>



Brian E. Whipker<sup>2</sup>

Volume 1 Number 12 March 2018

## Nutritional Monitoring Series

### Primula

(*Primula vulgaris*)

Primula require low fertility of 100 to 150 ppm N. They prefer a pH within the range of 5.5 to 6.2. Substrate pH values above 6.2 can inhibit iron availability and induce interveinal chlorosis or yellowing of the upper foliage. Primula are sensitive to excessive soluble salts [referred to as electrical conductivity (EC)] and may result in marginal leaf burn or necrosis.



Funding Generations of Progress  
Through Research and Scholarships



Project Sponsor



© Brian E. Whipker

Figure 1. High substrate pH above 6.2 will inhibit iron (Fe) uptake causing newly developed and recently matured leaves to become Fe-deficient and exhibit marginal and interveinal chlorosis (yellowing). Photos by: Brian E. Whipker

### Target Nutrition Parameters

**pH Category II,III:**

5.5 to 6.2

**Fertility Category:**

Low

100 to 150 ppm N

**EC Category A:**

1:2 Extraction:

0.4 to 0.6 mS/cm

SME:

0.9 to 1.3 mS/cm

PourThru:

1.3 to 2.0 mS/cm

Primula

[www.fertdirtandsquirt.com](http://www.fertdirtandsquirt.com)

<sup>1</sup>Michigan State University  
wgowen@msu.edu

<sup>2</sup>NC State University  
bwhipker@ncsu.edu



Figure 2. Substrate pH below 5.5 will inhibit magnesium (Mg) uptake causing lower or older leaves to become Mg-deficient and exhibit interveinal chlorosis. Photos by: Brian E. Whipker



Figure 3. High substrate pH above 6.2 will inhibit iron (Fe) uptake causing newly developed and recently matured leaves to become Fe-deficient and exhibit interveinal chlorosis (yellowing). Photo by: Brian E. Whipker

## Fertility Management of Primula

Micronutrient deficiencies or toxicities are common during primula production. Interveinal or marginal chlorosis (yellowing) or necrosis (death) may develop due to high substrate pH or over fertilization or high soluble salts [referred to as electrical conductivity (EC)].

Primula should be grown with a pH range of 5.5 to 6.2. Tissue nutrient levels found in healthy, newly expanded leaves and critical tissue values of primula are listed in Table 1. This range will enable growers to avoid high and low pH nutritional disorders.

Substrate pH above 6.2 will inhibit iron (Fe) uptake, causing newly developed and recently matured leaves to become Fe-deficient and exhibit marginal and interveinal chlorosis (Fig. 1). Corrective procedures for high substrate pH should begin within the range of 6.2 to 6.4. Maintaining substrate pH below 6.2 will reduce the likelihood of Fe-deficiency from occurring; however, applications of chelated micronutrients, such as Fe may be used to correct these deficiency symptoms.

Substrate pH below 5.5 will inhibit magnesium (Mg) uptake causing lower or older leaves to become Mg-deficient and exhibit interveinal chlorosis (Fig. 2) and intensify to overall leaf chlorosis (Fig. 3). Corrective procedures for low substrate pH should begin within the range of 5.3 to 5.5. Monthly applications of supplemental Mg in the form of magnesium sulfate ( $MgSO_4$ ; Epsom salts) at a rate of 8 oz./100 gallons of water in areas with naturally occurring Mg in the water supply or 16 oz./100 gallons of water in areas lacking Mg in the irrigation water (Whipker, personal communications) may be required if plants exhibit interveinal chlorosis of lower or older leaves.




Figure 4. Primula are sensitive to excessive soluble salts [referred to as electrical conductivity (EC)] and may result in marginal leaf burn or necrosis (death). Photo by: W. Garrett Owen

Table 1. Leaf tissue nutrient sufficiency range recommended for primula (*Primula vulgaris*).

Element		Sufficiency range <sup>1</sup>
Nitrogen (N)	(%)	2.50 - 3.30
Phosphorus (P)		0.40 - 0.80
Potassium (K)		1.80 - 3.00
Calcium (Ca)		0.60 - 1.00
Magnesium (Mg)		0.20 - 0.40
Sulfur (S)		0.18 - 0.29
Iron (Fe)	(ppm)	75 - 155
Manganese (Mn)		50 - 80
Zinc (Zn)		40 - 45
Copper (Cu)		5 - 10
Boron (B)		30 - 35
Molybdenum (Mo)		0.25 - 0.45

<sup>1</sup>Source: Bryson et al. (2014)

Plants may become chlorotic if substrate is kept too wet or drain poorly. Allow substrate to dry thoroughly between irrigations, but avoid permanent wilting.

Primula are highly sensitive to excessive EC levels. Substrate EC should remain below 0.6, 1.3, or 2.0 mS/cm, based on the 1:2 Extraction, SME, or PourThru methods, respectively. Accumulation of fertilizer salts in the substrate may result in marginal leaf burn or necrosis (Fig. 4) and mimic water stress symptomology. To avoid high EC, it is recommended to keep the fertilization rate low (100 to 150 ppm N). If levels begin to increase, switch to clear water irrigations to avoid a build-up of fertilizer salts in the substrate. If EC levels become excessive, then leach the substrate with clear irrigation water twice before providing fertility. It is best to monitor the crop to avoid excessive EC levels than to waste fertilizer by having to leach it out of the pots.

During propagation, specifically Stage 2 [radical, hypocotyl (stem), and cotyledons (seedling leaves) emergence] of seedling production, begin fertilizing at 50 to 75 ppm N provided from 14-0-14 (Dole and Wilkins, 2005; Karlsson, 1997). At Stage 3 (true leaves develop) and during production increase fertility to 100 to 150 ppm N provided from a well-balanced calcium nitrate [Ca(NO<sub>3</sub>)<sub>2</sub>] -based fertilizer with micronutrient supplements.

During winter months when plants are grown under cooler temperatures, fertilize with nitrate based fertilizers such as Ca(NO<sub>3</sub>)<sub>2</sub>. Avoid fertilizers providing ammonium (NH<sub>4</sub>-N) such as 20-10-20 or 20-20-20. Avoid ammoniacal-based fertilizers which can cause excessive vegetative growth and leaf expansion. Additionally, cold growing temperature or low fertility can result in premature bud set on small plants (Dole and Wilkins, 2005).

Primula can become phosphorous (P) deficient during production. Symptomology of P-deficient primula include newly developing leaves to curl inward and lower leaves become bronzed with brown veins (Karlsson, 1997).

### Summary

Providing primula with low fertility ranging from 100 to 150 ppm N, limit ammoniacal-based fertilizers, and maintaining a substrate pH of 5.5 to 6.2 will prevent most nutritional disorders.

### Literature Cited

Bryson, G.M., H.A. Mills, D.N. Sasseville, J.B. Jones, Jr., and A.V. Barker. 2014. Plant analysis handbook III. Micro-Macro Publishing, Inc., Athens, GA.

Dole, J.M. and H.F. Wilkins. 2005. Floriculture: Principles and species. 2nd ed. Pearson Education, Inc., Upper Saddle River, NJ.

Karlsson, M.G. 1997. Primula, p. 107-111. In: M.L. Gaston, S.A. Carver, C.A. Irwin, and R.A. Larson (eds.). Tips on growing specialty potted crops. Ohio Florist Association, Columbus, OH.



Funding Generations of Progress  
Through Research and Scholarships



Project Sponsor

## Corrective Procedures for Modifying Substrate pH and Electrical Conductivity (EC)

When the pH or substrate electrical conductivity (EC) drifts into unwanted territory, adjustments must be made. Below are the standard corrective procedures used to modify the substrate pH and EC for greenhouse grown crops in soilless substrates.

### 1. Low Substrate pH Correction

When Fe and Mn toxicity becomes a problem, adjust (raising) substrate pH to the recommended pH range. Corrective procedures to raise low pH levels are listed below. Switching to a basic fertilizer when the substrate pH is nearing the lower limit will help stabilize the pH. If the pH is below the recommended range, then corrective procedures will need to be implemented. Flowable lime is one option. Using a rate of 2 quarts per 100 gallons of water will typically increase the substrate pH by roughly 0.5 pH units. Two quarts can be used through an injector. Additional applications can be made if needed. Potassium bicarbonate ( $\text{KHCO}_3$ ) can also be applied. A rate of 2 pounds per 100 gallons of water will increase the substrate pH by roughly 0.8 pH units. This treatment will also provide excessive potassium (K) and cause a spike in the substrate EC. A leaching irrigation with clear water is required the following day to restore the nutrient balance (the ratio of K:Ca:Mg) and lower the EC. As always, remember to recheck your substrate pH to determine if reapplications are needed.

## pH Adjustment Recommendations

### *Flowable Lime*

- Use 1 to 2 quarts per 100 gallons of water.  
Rinse foliage.
- Avoid damage to your injector by using rates of 2 quarts per 100 gallons of water, or less.
- Can split applications.

### *Hydrated Lime*

- Mix 1 pound in 3 to 5 gallons of WARM water. Mix twice. Let settle. Decant liquid and apply through injector at 1:15.
- Caustic (rinse foliage ASAP and avoid skin contact)

### *Potassium Bicarbonate ( $\text{KHCO}_3$ )*

- Use 2 pounds per 100 gallons of water
- Rinse foliage immediately.
- Provides 933 ppm K.
- Leach heavily the following day with a complete fertilizer to reduce substrate EC and restore nutrient balance.
- Rates greater than 2 pounds per 100 gallons of water can cause phytotoxicity!

### 2. High Substrate pH Correction

The target pH for many species is between 5.8 and 6.2. Higher pH values will result in Fe deficiency and lead to the development of interveinal chlorosis on the upper leaves. Check the substrate pH to determine if it is too high. Be careful when lowering the substrate pH, because going too low can be much more problematic and difficult to deal with.

### *Acid-based Fertilizer*

If the substrate pH is just beginning to increase, then first consider switching to an acidic-based fertilizer. These ammoniacal-nitrogen (N) based fertilizers are naturally acidic and plant nitrogen uptake will help moderate the substrate pH over a week or two.

### *Acid Water Drench*

Some growers use this intermediate correction if pH levels are not excessively high and a quick lower of the substrate pH is desired. Use sulfuric acid to acidify your irrigation water to a pH 4.0 to 4.5. Apply this acid water as a substrate drench providing 5 to 10% excessive leaching of the substrate. Rinse the foliage to avoid phytotoxicity. Results should be visible within 5 days. Retest the substrate pH and repeat if needed.

### *Iron Drench*

If the levels are excessively high, then an Fe chelate application can be made to the substrate.

Below are the options.

### *Iron Chelate Drench (options)*

- Iron-EDDHA: mix 5 ounces in 100 gallons of water
- Iron-DTPA: mix 5 ounces in 100 gallons of water
- Iron sulfate: mix 4-8 ounces in 100 gallons of water
- Apply as a substrate drench with sufficient volume to leach the pot.
- Rinse foliage immediately.
- Avoid use on iron efficient plants (geraniums).

## 3. Low EC Correction

If low EC problems occur, increase the fertilization rate to 300 ppm N for a few applications before returning to the recommend fertilization rate for the crop.

## 4. High EC Correction

Excessively high fertilization rates will result in a marginal leaf burn. Check the substrate EC to confirm your diagnosis. Values greater than 6.0 mS/cm based on the PourThru sampling method can be problematic for many plants.

### *Switch to Clear Water Irrigations*

If the substrate EC is just beginning to increase over time, then leach with a few clear water irrigations to lower EC levels by flushing out the salts.

### *Clear Water Leaching*

If the EC values are excessively high, leach the substrate twice with back-to-back clear water irrigations. Then allow the substrate to dry down normally before retesting the EC. If EC levels are still too high, repeat the double leach. Once the substrate EC is back within the normal range, use a balanced fertilizer at a rate of 150 to 200 ppm N.



**e-GRO Alert**

[www.e-gro.org](http://www.e-gro.org)

**CONTRIBUTORS**

**Dr. Nora Catlin**  
Floriculture Specialist  
Cornell Cooperative Extension  
Suffolk County  
[nora.catlin@cornell.edu](mailto:nora.catlin@cornell.edu)

**Dr. Chris Currey**  
Assistant Professor of Floriculture  
Iowa State University  
[ccurrey@iastate.edu](mailto:ccurrey@iastate.edu)

**Dr. Ryan Dickson**  
Extension Specialist for Greenhouse  
Management & Technologies  
University of New Hampshire  
[ryan.dickson@unh.edu](mailto:ryan.dickson@unh.edu)

**Thomas Ford**  
Commercial Horticulture Educator  
Penn State Extension  
[tf2@psu.edu](mailto:tf2@psu.edu)

**Dan Gilrein**  
Entomology Specialist  
Cornell Cooperative Extension  
Suffolk County  
[dog1@cornell.edu](mailto:dog1@cornell.edu)

**Dr. Joyce Latimer**  
Floriculture Extension & Research  
Virginia Tech  
[jlatime@vt.edu](mailto:jlatime@vt.edu)

**Heidi Lindberg**  
Floriculture Extension Educator  
Michigan State University  
[wolleage@anr.msu.edu](mailto:wolleage@anr.msu.edu)

**Dr. Roberto Lopez**  
Floriculture Extension & Research  
Michigan State University  
[rglopez@msu.edu](mailto:rglopez@msu.edu)

**Dr. Neil Mattson**  
Greenhouse Research & Extension  
Cornell University  
[neil.mattson@cornell.edu](mailto:neil.mattson@cornell.edu)

**Dr. W. Garrett Owen**  
Floriculture Outreach Specialist  
Michigan State University  
[wgowen@msu.edu](mailto:wgowen@msu.edu)

**Dr. Rosa E. Raudales**  
Greenhouse Extension Specialist  
University of Connecticut  
[rosa.raudales@uconn.edu](mailto:rosa.raudales@uconn.edu)

**Dr. Beth Scheckelhoff**  
Extension Educator - Greenhouse Systems  
The Ohio State University  
[scheckelhoff.11@osu.edu](mailto:scheckelhoff.11@osu.edu)

**Lee Stivers**  
Extension Educator - Horticulture  
Penn State Extension  
Washington County  
[ljs32@psu.edu](mailto:ljs32@psu.edu)

**Dr. Paul Thomas**  
Floriculture Extension & Research  
University of Georgia  
[pathomas@uga.edu](mailto:pathomas@uga.edu)


**Dr. Ariana Torres-Bravo**  
Horticulture/ Ag. Economics  
Purdue University  
[torres2@purdue.edu](mailto:torres2@purdue.edu)

**Dr. Brian Whipker**  
Floriculture Extension & Research  
NC State University  
[bwhipker@ncsu.edu](mailto:bwhipker@ncsu.edu)

Copyright ©2018

Where trade names, proprietary products, or specific equipment are listed, no discrimination is intended and no endorsement, guarantee or warranty is implied by the authors, universities or associations.

**Cooperating Universities**



Cornell University

**Cornell Cooperative Extension  
Suffolk County**



**University of  
New Hampshire**  
Cooperative Extension

**PENNSSTATE**



Cooperative Extension  
College of Agricultural Sciences



**VIRGINIA  
TECH**

**MICHIGAN STATE  
UNIVERSITY**

**UCONN**

**PURDUE  
UNIVERSITY**



**The University of Georgia**



**THE OHIO STATE  
UNIVERSITY**

**NC STATE  
UNIVERSITY**

**IOWA STATE UNIVERSITY**

**In cooperation with our local and state greenhouse organizations**

**MAUMEE VALLEY GROWERS**  
*Choose the Very Best.*



Metro Detroit Flower Growers Association

